

Hyperspectral imaging data combined with climate data to predict stomatal conductance and transpiration of grapevine plants


Using Sequentially-Orthogonalized Partial-Least-Square Regression (SO-PLS)

Maxime Ryckewaert, Daphné Héran, Thierry Simonneau, Nicolas Saurin, Florent Abdelghafour, Romain Boulord, Daniel Moura, Silvia Mas-Garcia, Ryad Bendoula

INTRODUCTION

Digital agriculture driven by new intelligent sensors is one of the main ways to improve farm management.

Accessing physiological variables such as transpiration (E) and stomatal conductance (gs) in real time with optical instruments is challenging. These are the privileged variables to detect water stress.



OBJECTIVE

The objective is to evaluate predictions of transpiration (E) and stomatal conductance (gs) of grapevine (*Vitis vinifera* L.) by combining visible-near infrared spectral images with climate data, using Sequentially-Orthogonalized Partial-Least-Square Regression (SO-PLS).

EXPERIMENT

Pots of three grape varieties (Syrah, Merlot, Riesling) tested under two water conditions were studied to obtain water stress gradient. Hyperspectral images were acquired and a weather station provided radiation (Rg), relative humidity (RH), temperature (Ta) and wind speed (Ws). Precise monitoring of physiological variables was performed to obtain reference values.

Experimental design (Repeated twice)

Period: Summer 2020 / Location: Montpellier, la Gaillarde

Treatment	Varieties	Pots
Irrigated	Syrah	
None	Merlot	
	Riesling	

Hyperspectral and climate data acquisitions




Block 1

Hyperspectral imaging
Specim IQ
204 bands
between 397 and 1003 nm

Block 2

Temperature
Humidity
Global radiation
Wind Speed

Physiological variables



METHOD


Sequentially-Orthogonalized Partial-Least-Square Regression (SO-PLS) was used to build a prediction model.

$$y = Xb + Zc + r_{X,Z}$$


Steps of the SO-PLS

- 1 - fit y to X..... $y = Xb + r_X$
- 2 - orthogonalize Z to X-scores..... $Z = Z - T_X (T_X^T T_X)^{-1} T_X^T Z$
- 3 - fit rx to Z $r_X = Z c + r_{X,Z}$

RESULTS




MODEL EVALUATION



- A calibration set of 106 hyperspectral images
- A test set of 54 hyperspectral images (reflecting all modalities of the experimental design)

MODEL APPLICATION



- Opportunity to study spatial distribution of stresses over the leaf stages
- Within-variability as a new phenotyping trait for breeding varieties

CONCLUSION AND PERSPECTIVES

The combination of climate data and hyperspectral images provides models with good performance to predict physiological variables. Quality of these prediction models could be improved by defining varietal models on a larger data set. These encouraging results offer prospects for the use of spectral imaging to study water stress of grapevine plants.



Reference

Ryckewaert, Maxime, Daphné Héran, Thierry Simonneau, Florent Abdelghafour, Romain Boulord, Nicolas Saurin, Daniel Moura, Silvia Mas-Garcia, et Ryad Bendoula. « Physiological Variable Predictions Using VIS-NIR Spectroscopy for Water Stress Detection on Grapevine: Interest in Combining Climate Data Using Multiblock Method ». Computers and Electronics in Agriculture 197 (juin 2022): 106973. <https://doi.org/10.1016/j.compag.2022.106973>.



Contact:

maxime.ryckewaert@inrae.fr
ryad.bendoula@inrae.fr